

Resonant Bound State Production at e^-e^- Colliders

David Bowser-Chao¹, Tom D. Imbo², B. Alex King³ and Eric C. Martell⁴

Department of Physics
University of Illinois at Chicago
845 W. Taylor St.
Chicago, IL 60607-7049

Observation of a sequence of resonances at an e^-e^- collider would suggest bound states of strongly coupled constituents carrying lepton number. Obvious candidates for these exotic constituents are leptoquarks and leptogluons. We show that under reasonable assumptions, the existence of one leptogluon flavor of appropriate mass can give rise to sizeable “leptoglueball” production rates and observable resonance peaks. In contrast, one needs two leptoquark flavors in order to produce the analogous “leptoquarkonium” states. Moreover, cross-generational leptoquark couplings are necessary to give observable event rates in many cases, and leptoquarkonium mass splittings are too small to resolve with realistic beam energy resolutions.

¹ davechao@hepster.phy.uic.edu

² imbo@uic.edu

³ aking@uic.edu

⁴ ecm@uic.edu

Numerous TeV-scale colliders have been proposed as possible complements to the planned Large Hadron Collider (LHC). For example, e^+e^- , $\mu^+\mu^-$, $e^-\gamma$, $\gamma\gamma$ and e^-e^- beams are all being seriously considered. Because of their lepton number $L = 2$ initial state, e^-e^- colliders are generally thought to have the least physics potential. However such a facility could, for example, study strong W^-W^- scattering — an aspect of electroweak symmetry breaking not readily accessible at the LHC or other machines [1]. Moreover, if doubly-charged particles with $L = 2$ exist and have reasonable couplings to electrons, they would be produced copiously at an e^-e^- collider through an s -channel resonance [2][3]. If such *bileptons* are fundamental (or have a compositeness scale which is well above the collider energy range), then there will be a single resonance peak. If, however, these bileptons are composite at these energy scales, then there should be a sequence of peaks representing excitations of the underlying constituents, allowing a spectroscopic study similar to well-known treatments of quarkonium systems.

Because they carry lepton number and have strong interactions, *colored leptons* provide a possible mechanism for the formation of such bound states. For example, *leptogluons* (LG s) can occur in quark/lepton composite models containing colored preons [4][5][6]. LG s would be color octets, like Standard Model gluons, but would also carry lepton number, electric charge, and have a spin of 1/2 or 3/2. They would have all the couplings of ordinary leptons and gluons, as well as additional trilinear couplings to lepton-gluon pairs. Current experiments tell us that, if they exist, spin-1/2 LG s which couple to e^-g must have masses $m_{LG} \gtrsim 325$ GeV [6]. We will show that a single flavor of such LG will give rise, under reasonable assumptions, to sizeable production rates of LG - LG bound states (leptoglueballs) at an e^-e^- collider. Moreover, even given the poor beam energy resolution expected at such a machine, the bound state mass splittings are big enough (and widths small enough) so that some resonance peaks can likely be reconstructed.

The relevant (non-renormalizable) interaction term in the lagrangian for spin-1/2 LG s has the form [6]

$$\frac{g_s}{\Lambda} [\lambda_L \bar{e}_R \sigma^{\mu\nu} G_L^a + \lambda_R \bar{\ell}_L \sigma^{\mu\nu} G_R^a] F_{\mu\nu}^a + h.c. \quad . \quad (1)$$

Here $\bar{\ell}_L$ is the left-handed first generation $SU(2)_L$ lepton doublet, G_R^a the right-handed LG doublet, e_R and G_L^a the corresponding singlets, $F_{\mu\nu}^a$ the $SU(3)_c$ field strength tensor, and $g_s = \sqrt{4\pi\alpha_s(\mu)}$ the QCD running coupling. Defining the dimensionless couplings λ_L and λ_R to be of order 1, Λ represents the “compositeness scale” of the LG . In order to avoid possible problems with magnetic moment constraints, we assume a purely chiral

interaction. Specifically, we take $\lambda_R = 0$ and $\lambda_L = 1$ so that the leptogluon only couples to the right-handed electron e_R . (The choice $\lambda_R = 1$, $\lambda_L = 0$ gives equivalent results for the charge -1 component of the right-handed LG doublet.) If the leptogluon and electron are simply different color states of the same underlying preon configuration, then m_{LG} will be $\mathcal{O}(\Lambda\alpha_s(\Lambda))$ [4].

We now discuss leptoglueball production, restricting our attention to S -wave ($\ell = 0$) states. (The production of P -wave states is suppressed relative to these by a factor of order α_s^2 .) Since these leptoglueballs are formed from identical fermions, their overall wavefunctions must be antisymmetric under the simultaneous interchange of color, spin and spatial variables. Hence, for color singlet S -states, the two LG s must be in the antisymmetric spin-0 state. (This is analogous to the situation for bound states of two gluinos [7].) There are two primary decay channels for these leptoglueballs, namely the annihilation of the two LG s (which occurs via t -channel gluon exchange)

$$\Gamma_{B \rightarrow e_R^- e_R^-} = 288\pi \alpha_s(m_{LG})^2 |\psi(0)|^2 \frac{M_B^2}{\Lambda^4}, \quad (2)$$

and the spectator decay of one of the LG s

$$\Gamma_{B \rightarrow e_R^- g + X} = 2\Gamma_{LG \rightarrow e_R^- g} = 2\alpha_s(m_{LG}) \frac{m_{LG}^3}{\Lambda^2}. \quad (3)$$

Here M_B is the mass of the nonrelativistic bound state B and $|\psi(0)|^2$ the square of its wavefunction at the origin. Since these LG s must be very heavy, it is reasonably accurate to describe leptoglueballs using the perturbative Coulombic potential $V(r) = -\frac{C_2\alpha_s(\mu)}{r}$. (We will ignore hyperfine effects.) For the nS state this leads to $M_B \simeq 2m_{LG} - \left(\frac{C_2\alpha_s(\mu_n)}{2n}\right)^2 m_{LG}$ and $|\psi(0)|^2 \simeq \frac{1}{\pi} \left(\frac{C_2\alpha_s(\mu_n)}{2n} m_{LG}\right)^3$. The ‘‘Casimir factor’’ C_2 is 3 for color octets (compared to $4/3$ for color triplets), and $\mu_n = \frac{C_2\alpha_s(m_{LG})}{2n} m_{LG}$ is a typical momentum scale for the nS state. For all channels this gives $\Gamma_{B \rightarrow e_R^- e_R^-} \ll \Gamma_{B \rightarrow e_R^- g + X}$, and the total width is well-approximated as $\Gamma(B) \simeq \Gamma_{B \rightarrow e_R^- g + X}$.

Using the above results, we can now compute the total production cross section, σ_{tot} , for $e_R^- e_R^- \rightarrow B$:

$$\sigma_{\text{tot}} = \frac{32\pi}{M_B^2} \frac{\Gamma_{B \rightarrow e_R^- e_R^-}}{\Gamma(B)} \cdot f(x), \quad (4)$$

where $f(x) = 2x e^{x^2} \int_x^\infty e^{-t^2} dt$ and $x = \frac{1}{\sqrt{8}} \frac{\Gamma(B)}{\sigma_E}$. The center-of-mass energy has been assumed to have a gaussian distribution peaked at the resonance mass M_B with a spread of

σ_E . The factor f takes into account the effects of this beam energy uncertainty: $f(x) \rightarrow 1$ as $\sigma_E \rightarrow 0$ and $f(x) \simeq \sqrt{\pi} x$ for $\sigma_E \gg \Gamma(B)$. For nS states in the above Coulombic approximation, (4) becomes simply

$$\sigma_{\text{tot}} = 15552\pi \cdot \frac{\alpha_s(m_{LG})^3 \alpha_s(\mu_n)^3}{m_{LG}^2 n^3} \cdot f(x), \quad (5)$$

where we have set $\Lambda = \frac{m_{LG}}{\alpha_s(m_{LG})}$. The expected beam energy resolution for a TeV-scale e^-e^- collider is $\sigma_E \simeq 0.01\sqrt{s} \gtrsim 10\Gamma(B)$, giving an $f(x)$ on the order of 0.05.

We have used (5) to calculate the expected number of events for the production of $1S$ leptogluoballs on resonance (see Table 1). The event rates for the $2S$ (respectively, $3S$) states will be smaller by about a factor of 6 (respectively, 16). Note that these calculations did not require any unnatural assumptions about the strength of the coupling to produce sizeable rates. These rates should be several orders of magnitude larger than the QED background $e^-e^- \rightarrow e^-e^- + (\gamma^* \rightarrow 2 \text{ jets})$. In addition, since a leptogluoball (at rest) decaying into $e^-e^- + 2 \text{ jets}$ will look very different from this QED process, the application of appropriate cuts should reduce this background even further. In the Coulombic approximation, the mass splitting between the $1S$ and $2S$ states (Δ_{2S-1S}) ranges from ~ 10 to 20 GeV as m_{LG} varies from 400 to 1000 GeV . This is about 14 times bigger than the natural width $\Gamma(B)$ of the states for all values of m_{LG} . (Δ_{3S-2S} is about $3\Gamma(B)$.) Moreover, the distance between the $1S$ resonance peak and the continuum LG pair production threshold at $\sqrt{s} = 2m_{LG}$ is $\sim 20\Gamma(B)$. Thus if we had unlimited beam resolution, at least the $1S$ and $2S$ resonance peaks should be easily visible. However, the expected beam energy spread σ_E is about the same as Δ_{2S-1S} . Even so, it may still be possible to reconstruct the $1S$ state, as well as some evidence for the $2S$ state, from a detailed analysis of the shape of the observed cross section. Obtaining a σ_E which is a factor of 5 to 10 smaller than expected would not only go a long way in helping us to see these successive peaks, but would also increase the production cross sections by increasing $f(x)$.¹

¹ One can also consider the production of “second generation” leptogluons at a $\mu^-\mu^-$ collider, which is expected to have a much better beam energy resolution of $\sim 2 \times 10^{-5}\sqrt{s}$ [8].

m_{LG}	$e_R^- e_R^- \rightarrow B$
400	1.472×10^5
600	4.267×10^4
800	1.796×10^4
1000	9.243×10^3

Table 1: Estimates of the number of $1S$ leptoglueball events on resonance at a polarized e^-e^- collider. The numbers are obtained from (5) using the three-loop formula for $\alpha_s(\mu)$ given in [9]. Masses are in GeV and the integrated luminosity is taken to be 10 fb^{-1} .

It is interesting to compare the above leptoglueballs with *fundamental* scalar bileptons S of the same mass and quantum numbers. If we denote the (dimensionless) trilinear coupling of S to two right-handed electrons by g , then we would like to know the “effective” g for leptoglueballs. That is, for what value of g does $\Gamma_{B \rightarrow e_R^- e_R^-}$ from (2) equal $\Gamma_{S \rightarrow e_R^- e_R^-} = \frac{g^2 M_S}{8\pi}$ [3]? For example, setting $M_S = M_B = 800 \text{ GeV}$ these two widths are equal (for $1S$ leptoglueballs) if $g \simeq \frac{1}{100}$, where again we have used the Coulombic approximation and $\Lambda = \frac{m_{LG}}{\alpha_s(m_{LG})}$. Thus, the effective coupling of leptoglueballs to electron pairs is small but non-negligible.

So far, the focus of this letter has been the resonant production of composite bileptons at e^-e^- colliders with spin-1/2 leptogluons as the constituents. Of course, we can also consider the production of these hypothetical LG s at other future colliders. For example, the best scenario for the *discovery* of LG s would most likely be single production at an e^-p machine. There are also lepton-number-zero analogs of our leptoglueballs, namely $LG-\overline{LG}$ bound states, which can in principle be produced on resonance at an e^+e^- collider. Ignoring extremely small electroweak effects, spin-zero $LG-\overline{LG}$ states with $\ell = 0$ will have the same spectrum as our $LG-LG$ states above, although with somewhat larger widths due to additional annihilation decay channels. However, the cross section for resonant production of these scalar bound states at an e^+e^- machine is proportional to $(\lambda_L \lambda_R)^2$, requiring both λ_L and λ_R in (1) to be non-zero. The measured electron magnetic moment then constrains $\lambda_L \lambda_R$ to be very small, resulting in a huge suppression. Thus, the spin-0 $LG-LG$ states that we have considered in the context of e^-e^- collisions have no observable $LG-\overline{LG}$ analog (on resonance) at an e^+e^- collider. It *will* be possible to produce spin-1 $LG-\overline{LG}$ bound states with $\ell = 0$ on resonance in e^+e^- collisions (with reasonably high

rates and low backgrounds). Interestingly, these states have no analog in LG - LG systems due to the exclusion principle. So we see that e^+e^- and e^-e^- colliders will allow us to study complementary aspects of the spectroscopy of bound states of leptogluons.

Finally, we will briefly consider another candidate for the constituents of composite bileptons, namely *leptoquarks* (LQ s) — color triplets (or antitriplets) carrying lepton number and having a spin of either 0 or 1. These objects have been the subject of much discussion over the last few years (see, for example, the papers cited in [10]). However the resonant production of a bound state of two LQ s (leptoquarkonia) at an e^-e^- collider is somewhat unnatural, requiring two distinct leptoquark flavors: one which couples to e^-q and one which couples to $e^-\bar{q}$ for some quark flavor q .² In addition, many such leptoquarkonium production rates are “quark mass suppressed” due to a necessary chirality flip of the exchanged t -channel quark. Examples of this are the production of an S -wave bound state of two scalar LQ s or of one scalar and one vector LQ . In such cases, a cross-generational coupling to the top quark is necessary to produce non-negligible event rates. The cross section for the production of S -wave bound states of two *vector* LQ s from incoming electrons with opposite helicity does *not* have the above quark mass suppression. (P -wave bound states can also be produced without a chirality flip regardless of the LQ spin.) But in any case, even if the above leptoquarkonium states do exist and have reasonable production rates, the weaker QCD interactions of color triplets versus color octets will lead to bound state mass splittings about 4 times smaller than those of leptogluoballs of similar mass. Since these splittings are much smaller than the expected beam energy resolution, it is unlikely that any resonance peaks could be reconstructed. All we are likely to see is a broad enhancement of the continuum LQ pair production threshold, with no obvious sign of a resonance peak.

In conclusion, we have seen that the existence of spin-1/2 leptogluons of appropriate mass which couple to e^-g could give rise to striking bound state resonances (leptogluoballs) at a TeV-scale e^-e^- collider. These bileptonic resonances would arise more naturally and be much easier to observe than those produced by leptoquarks of similar mass. The production of a pair of identical 1 TeV LG s will lead, under reasonable assumptions, to $1S$ and $2S$ scalar leptogluoball resonance peaks well above background, separated by ~ 20 GeV, each having a width of about 1.4 GeV. The $2S$ peak, which is $\sim 1/6$ the height

² Production of bound states containing leptoquarks at other future colliders has been discussed in [9]. For a discussion of continuum leptoquark production at e^-e^- colliders, see [11].

of the $1S$ peak, remains about 9 GeV below the continuum LG pair production threshold. One hopes that a sufficiently small beam resolution can be achieved in order to allow us to resolve these peaks. This analysis is sensitive only to the assumption that $\Lambda = \frac{m_{LG}}{\alpha_s(m_{LG})}$. Although this seems natural from a composite model point of view, (2)–(4) tell us that any significant deviation from this equality would greatly affect both $\Gamma(B)$ and σ_{tot} (while not changing the bound state mass splittings).

Acknowledgements

It is a pleasure to acknowledge Lee Brekke, Adam Falk, Howard Georgi, Tao Han, Wai-Yee Keung, Young-Kee Kim, Joe Lykken, Uday Sukhatme, Scott Willenbrock and Dieter Zeppenfeld for helpful discussions. A very special thank you goes to Phil Jackson for years of guidance. This research was supported in part by the U.S. Department of Energy under Grant Number DE-FG02-91ER40676.

References

- [1] H. Murayama, Int. J. Mod. Phys. A **13** (1998) 2227;
T. Han, Int. J. Mod. Phys. A **13** (1998) 2337.
- [2] P. Frampton, Int. J. Mod. Phys. A **13** (1998) 2345.
- [3] M. Raidal, Phys. Rev. D **57** (1998) 2013;
F. Cuypers and M. Raidal, Nucl. Phys. B **501** (1997) 3.
- [4] H. Fritzsch and G. Mandelbaum, Phys. Lett. **102B** (1981) 319;
H. Harari and N. Seiberg, Phys. Lett. **98B** (1981) 269;
E. J. Squires, Phys. Lett. **94B** (1980) 54.
- [5] T. G. Rizzo, Phys. Rev. D **34** (1986) 133; Phys. Rev. D **33** (1986) 1852;
Y. Nir, Phys. Lett. **164B** (1986) 395;
K. H. Streng, Z. Phys. C **33** (1986) 247;
U. Baur and K. H. Streng, Z. Phys. C **30** (1986) 325; Phys. Lett. **162B** (1985) 387;
H. Harari, Phys. Lett. **156B** (1985) 250;
S. F. King and S. R. Sharpe, Nucl. Phys. B **253** (1985) 1; Phys. Lett. **143B** (1984) 494.
- [6] J. L. Hewitt and T. G. Rizzo, Phys. Rev. D **56** (1997) 5709.
- [7] W.-Y. Keung and A. Khare, Phys. Rev. D **29** (1984) 2657.
- [8] M. Demarteau and T. Han, hep-ph/9801407.
- [9] D. Bowser-Chao, T. D. Imbo, B. Alex King and E. C. Martell, Phys. Lett. B **432** (1998) 167.
- [10] P. H. Frampton and M. Harada, hep-ph/9711448.
- [11] F. Cuypers, P. Frampton and R. Rückl, Phys. Lett. B **390** (1997) 221.